

A probability inequality for sums of independent Banach space valued random variables

Deli Li¹, Han-Ying Liang^{2,*}, and Andrew Rosalsky³

¹Department of Mathematical Sciences, Lakehead University,
Thunder Bay, Ontario, Canada

²Department of Mathematics, Tongji University,
Shanghai, China

³Department of Statistics, University of Florida,
Gainesville, Florida, USA

Abstract Let $(\mathbf{B}, \|\cdot\|)$ be a real separable Banach space. Let $\varphi(\cdot)$ and $\psi(\cdot)$ be two continuous and increasing functions defined on $[0, \infty)$ such that $\varphi(0) = \psi(0) = 0$, $\lim_{t \rightarrow \infty} \varphi(t) = \infty$, and $\frac{\psi(\cdot)}{\varphi(\cdot)}$ is a nondecreasing function on $[0, \infty)$. Let $\{V_n; n \geq 1\}$ be a sequence of independent and symmetric $\mathbf{B}$ -valued random variables. In this note, we establish a probability inequality for sums of independent $\mathbf{B}$ -valued random variables by showing that for every $n \geq 1$ and all $t \geq 0$,

$$\mathbb{P}\left(\left\|\sum_{i=1}^n V_i\right\| > tb_n\right) \leq 4\mathbb{P}\left(\left\|\sum_{i=1}^n \varphi(\psi^{-1}(\|V_i\|)) \frac{V_i}{\|V_i\|}\right\| > ta_n\right) + \sum_{i=1}^n \mathbb{P}(\|V_i\| > b_n),$$

where $a_n = \varphi(n)$ and $b_n = \psi(n)$, $n \geq 1$. As an application of this inequality, we establish what we call a comparison theorem for the weak law of large numbers for independent and identically distributed $\mathbf{B}$ -valued random variables.

Keywords Probability inequality · Sums of independent Banach space valued random variables · Weak law of large numbers

Mathematics Subject Classification (2000): 60E15 · 60B12 · 60G50

Running Head: Probability inequality

1 Introduction and the main result

We begin with some notation. Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and let $(\mathbf{B}, \|\cdot\|)$ be a real separable Banach space equipped with its Borel σ -algebra $\mathcal{B}$ (= the σ -algebra generated by the class of open subsets of $\mathbf{B}$ determined by $\|\cdot\|$). A $\mathbf{B}$ -valued random variable X is defined as a measurable function from $(\Omega, \mathcal{F})$ into $(\mathbf{B}, \mathcal{B})$. Throughout this note, let $\{R, R_n; n \geq 1\}$ be a Rademacher sequence; i.e., $\{R, R_n; n \geq 1\}$ is a sequence of independent and identically distributed (i.i.d.) real-valued random variables with $\mathbb{P}(R = -1) = \mathbb{P}(R = 1) = 1/2$.

Probability inequalities are essential for establishing virtually all probability limit theorems and for advancing statistical theory, and they also have an intrinsic interest as well. For example, some invaluable and celebrated classical inequalities are those of Lévy, Ottaviani, Kahane, Hoffmann-Jørgensen (see, e.g., Lévy (1937), Chow and Teicher (1997, p. 75), Kahane (1968), and Hoffmann-Jørgensen (1974), respectively), etc. This note is devoted to establishing the following probability inequality which is a comparison theorem for sums of independent $\mathbf{B}$ -valued random variables.

*Corresponding author: Han-Ying Liang (Telephone: 86-21-65983242, FAX: 86-21-65983242)

Theorem 1.1. Let $\varphi(\cdot)$ and $\psi(\cdot)$ be two continuous and increasing functions defined on $[0, \infty)$ such that $\varphi(0) = \psi(0) = 0$ and

$$\lim_{t \rightarrow \infty} \varphi(t) = \infty \quad \text{and} \quad \frac{\psi(\cdot)}{\varphi(\cdot)} \text{ is a nondecreasing function on } [0, \infty). \quad (1.1)$$

Here we define $\frac{\varphi(0)}{\psi(0)} = \lim_{t \rightarrow 0^+} \frac{\varphi(t)}{\psi(t)}$. For $n \geq 1$, set $a_n = \varphi(n)$ and $b_n = \psi(n)$. Then we have:

(i) Let $\{x_n; n \geq 1\}$ be a $\mathbf{B}$ -valued sequence such that $\|x_n\| \leq b_n$, $n \geq 1$. Then for every $n \geq 1$ and all $t \geq 0$,

$$\mathbb{P} \left(\left\| \sum_{i=1}^n R_i x_i \right\| > t b_n \right) \leq 2 \mathbb{P} \left(\left\| \sum_{i=1}^n R_i \varphi(\psi^{-1}(\|x_i\|)) \frac{x_i}{\|x_i\|} \right\| > t a_n \right). \quad (1.2)$$

Here $\varphi(\psi^{-1}(\|0\|)) \frac{0}{\|0\|} \triangleq 0$ since $\lim_{x \rightarrow 0} \varphi(\psi^{-1}(\|x\|)) \frac{x}{\|x\|} = 0$.

(ii) If $\{V_n; n \geq 1\}$ is a sequence of independent and symmetric $\mathbf{B}$ -valued random variables, then for every $n \geq 1$ and all $t \geq 0$,

$$\mathbb{P} \left(\left\| \sum_{i=1}^n V_i \right\| > t b_n \right) \leq 4 \mathbb{P} \left(\left\| \sum_{i=1}^n \varphi(\psi^{-1}(\|V_i\|)) \frac{V_i}{\|V_i\|} \right\| > t a_n \right) + \sum_{i=1}^n \mathbb{P}(\|V_i\| > b_n). \quad (1.3)$$

The proof of Theorem 1.1 is given in Section 2. An application of Theorem 1.1 is presented in Section 3. The application provides what we refer to as a comparison theorem for the weak law of large numbers (WLLN) for i.i.d. $\mathbf{B}$ -valued random variables.

2 Proof of Theorem 1.1

To establish Theorem 1.1, we use the following result which is the second part of Theorem 4.4 of Ledoux and Talagrand (1991).

Lemma 2.1. Let $\{x_n; n \geq 1\}$ be a $\mathbf{B}$ -valued sequence and let $\{\alpha_n; n \geq 1\}$ be a real-valued sequence such that $\sup_{n \geq 1} |\alpha_n| \leq 1$. Then we have, for every $n \geq 1$ and all $t \geq 0$,

$$\mathbb{P} \left(\left\| \sum_{i=1}^n \alpha_i R_i x_i \right\| > t \right) \leq 2 \mathbb{P} \left(\left\| \sum_{i=1}^n R_i x_i \right\| > t \right).$$

Proof of Theorem 1.1 Part (i). Clearly we have, for every $n \geq 1$ and all $t \geq 0$,

$$\begin{aligned} \mathbb{P} \left(\left\| \sum_{i=1}^n R_i x_i \right\| > t b_n \right) &= \mathbb{P} \left(\left\| \sum_{i=1}^n \left(\frac{a_n}{b_n} \cdot \frac{\psi(\psi^{-1}(\|x_i\|))}{\varphi(\psi^{-1}(\|x_i\|))} \right) R_i \frac{\varphi(\psi^{-1}(\|x_i\|))}{\psi(\psi^{-1}(\|x_i\|))} x_i \right\| > t a_n \right) \\ &= \mathbb{P} \left(\left\| \sum_{i=1}^n \left(\frac{\varphi(n)}{\psi(n)} \cdot \frac{\psi(\psi^{-1}(\|x_i\|))}{\varphi(\psi^{-1}(\|x_i\|))} \right) R_i \varphi(\psi^{-1}(\|x_i\|)) \frac{x_i}{\|x_i\|} \right\| > t a_n \right). \end{aligned}$$

Since $\varphi(\cdot)$ and $\psi(\cdot)$ are two continuous and increasing functions defined on $[0, \infty)$ satisfying $\varphi(0) = \psi(0) = 0$ and (1.1), we see that $\varphi^{-1}(\cdot)$ is also a continuous and increasing function defined on $[0, \infty)$ such that $\psi^{-1}(0) = 0$, $\lim_{t \rightarrow \infty} \psi^{-1}(t) = \infty$, and

$$0 \leq \frac{\psi(\psi^{-1}(t))}{\varphi(\psi^{-1}(t))} \leq \frac{\psi(n)}{\varphi(n)} = \frac{b_n}{a_n} \quad \text{whenever} \quad 0 \leq t \leq \psi(n) = b_n.$$

Note that $\|x_n\| \leq b_n = \psi(n)$, $n \geq 1$. We thus conclude that, for every $n \geq 1$,

$$0 \leq \frac{\psi(\psi^{-1}(\|x_i\|))}{\varphi(\psi^{-1}(\|x_i\|))} \leq \frac{\psi(n)}{\varphi(n)} \quad \text{for } i = 1, 2, \dots, n$$

and hence that, for every $n \geq 1$,

$$0 \leq \frac{\varphi(n)}{\psi(n)} \cdot \frac{\psi(\psi^{-1}(\|x_i\|))}{\varphi(\psi^{-1}(\|x_i\|))} \leq 1 \quad \text{for } i = 1, 2, \dots, n.$$

By applying Lemma 2.1 we thus have, for every $n \geq 1$ and all $t \geq 0$,

$$\begin{aligned} & \mathbb{P} \left(\left\| \sum_{i=1}^n \left(\frac{\varphi(n)}{\psi(n)} \cdot \frac{\psi(\psi^{-1}(\|x_i\|))}{\varphi(\psi^{-1}(\|x_i\|))} \right) R_i \varphi(\psi^{-1}(\|x_i\|)) \frac{x_i}{\|x_i\|} \right\| > ta_n \right) \\ & \leq 2\mathbb{P} \left(\left\| \sum_{i=1}^n R_i \varphi(\psi^{-1}(\|x_i\|)) \frac{x_i}{\|x_i\|} \right\| > ta_n \right) \end{aligned}$$

proving Part **(i)**.

Part **(ii)**. For every $n \geq 1$, write

$$V_{n,i} = V_i I \{ \|V_i\| \leq b_n \}, \quad T_i = \varphi(\psi^{-1}(\|V_i\|)) \frac{V_i}{\|V_i\|}, \quad T_{n,i} = \varphi(\psi^{-1}(\|V_{n,i}\|)) \frac{V_{n,i}}{\|V_{n,i}\|}, \quad i = 1, \dots, n.$$

Clearly we have, for all $t \geq 0$,

$$\mathbb{P} \left(\left\| \sum_{i=1}^n V_i \right\| > tb_n \right) \leq \mathbb{P} \left(\left\| \sum_{i=1}^n V_{n,i} \right\| > tb_n \right) + \sum_{i=1}^n \mathbb{P}(\|V_i\| > b_n). \quad (2.1)$$

Note that

$$\{\|V_i\| \leq b_n\} = \{\psi^{-1}(\|V_i\|) \leq n\} = \{\varphi(\psi^{-1}(\|V_i\|)) \leq a_n\} = \{\|T_i\| \leq a_n\}, \quad i = 1, \dots, n.$$

Thus it is easy to see that

$$T_{n,i} = T_i I \{ \|T_i\| \leq a_n \}, \quad i = 1, \dots, n.$$

Since $\{V_i; i \geq 1\}$ is a sequence of independent and symmetric $\mathbf{B}$ -valued random variables, $\{V_{n,i}; i = 1, \dots, n\}$, $\{T_i; i = 1, \dots, n\}$, and $\{T_{n,i}; i = 1, \dots, n\}$ are finite sequences of independent and symmetric $\mathbf{B}$ -valued random variables. Let $\{R, R_n; n \geq 1\}$ be a Rademacher sequence which is independent of $\{V_n; n \geq 1\}$. Then $\{R_i V_{n,i}; i = 1, \dots, n\}$ has the same distribution as $\{V_{n,i}; i = 1, \dots, n\}$ in $\mathbf{B}^n$ and $\{R_i T_{n,i}; i = 1, \dots, n\}$ has the same distribution as $\{T_{n,i}; i = 1, \dots, n\}$ in $\mathbf{B}^n$. Since $\|V_{n,i}\| \leq$

b_n , $i = 1, \dots, n$, by applying (1.2), we have, for all $t \geq 0$,

$$\begin{aligned}
\mathbb{P} \left(\left\| \sum_{i=1}^n V_{n,i} \right\| > tb_n \right) &= \mathbb{P} \left(\left\| \sum_{i=1}^n R_i V_{n,i} \right\| > tb_n \right) \\
&= \mathbb{E} \left(\mathbb{P} \left(\left\| \sum_{i=1}^n R_i V_{n,i} \right\| > tb_n \middle| V_1, \dots, V_n \right) \right) \\
&\leq 2\mathbb{E} \left(\mathbb{P} \left(\left\| \sum_{i=1}^n R_i T_{n,i} \right\| > ta_n \middle| V_1, \dots, V_n \right) \right) \quad (2.2) \\
&= 2\mathbb{P} \left(\left\| \sum_{i=1}^n R_i T_{n,i} \right\| > ta_n \right) \\
&= 2\mathbb{P} \left(\left\| \sum_{i=1}^n T_{n,i} \right\| > ta_n \right).
\end{aligned}$$

For all $n \geq 1$, since $\{T_i; i = 1, \dots, n\}$ is a finite sequence of independent and symmetric $\mathbf{B}$ -valued random variables, it follows that $\{T_i I \{\|T_i\| \leq a_n\} - T_i I \{\|T_i\| > a_n\}; i = 1, \dots, n\}$ has the same distribution as $\{T_i; i = 1, \dots, n\}$ in $\mathbf{B}^n$. Note that

$$\sum_{i=1}^n T_{n,i} = \frac{\sum_{i=1}^n T_i + \sum_{i=1}^n (T_i I \{\|T_i\| \leq a_n\} - T_i I \{\|T_i\| > a_n\})}{2}, \quad n \geq 1.$$

We thus have, for every $n \geq 1$ and all $t \geq 0$,

$$\begin{aligned}
\mathbb{P} \left(\left\| \sum_{i=1}^n T_{n,i} \right\| > ta_n \right) &\leq \mathbb{P} \left(\left\| \sum_{i=1}^n T_i \right\| > ta_n \right) \\
&\quad + \mathbb{P} \left(\left\| \sum_{i=1}^n (T_i I \{\|T_i\| \leq a_n\} - T_i I \{\|T_i\| > a_n\}) \right\| > ta_n \right) \quad (2.3) \\
&= 2\mathbb{P} \left(\left\| \sum_{i=1}^n T_i \right\| > ta_n \right).
\end{aligned}$$

Now we can see that (1.3) follows from (2.1), (2.2), and (2.3). $\square$

3 An application

As an application of Theorem 1.1, in this section we will establish what we call a comparison theorem for the WLLN for i.i.d. $\mathbf{B}$ -valued random variables. Theorem 3.1 is new even when $\mathbf{B} = \mathbb{R}$.

Theorem 3.1. (Comparison theorem for the WLLN). *Let $(\mathbf{B}, \|\cdot\|)$ be a real separable Banach space. Let $\{a_n; n \geq 1\}$ and $\{b_n; n \geq 1\}$ be increasing sequences of positive real numbers such that*

$$\lim_{n \rightarrow \infty} a_n = \infty \text{ and } \{b_n/a_n; n \geq 1\} \text{ is a nondecreasing sequence.} \quad (3.1)$$

Suppose that, for every symmetric sequence $\{X, X_n; n \geq 1\}$ of i.i.d. $\mathbf{B}$ -valued random variables,

$$\frac{S_n}{a_n} \rightarrow_{\mathbb{P}} 0 \text{ if and only if } \lim_{n \rightarrow \infty} n\mathbb{P}(\|X\| > a_n) = 0. \quad (3.2)$$

Here and below $S_n = \sum_{i=1}^n X_i$, $n \geq 1$. Then, for every sequence $\{X, X_n; n \geq 1\}$ of i.i.d. $\mathbf{B}$ -valued random variables, we have that

$$\frac{S_n - \gamma_n}{b_n} \rightarrow_{\mathbb{P}} 0 \text{ or } \limsup_{n \rightarrow \infty} \mathbb{P}\left(\frac{\|S_n - \gamma_n\|}{b_n} > \lambda\right) > 0 \quad \forall \lambda > 0 \quad (3.3)$$

according as

$$\lim_{n \rightarrow \infty} n\mathbb{P}(\|X\| > b_n) = 0 \text{ or } \limsup_{n \rightarrow \infty} n\mathbb{P}(\|X\| > b_n) > 0. \quad (3.4)$$

Here and below $\gamma_n = n\mathbb{E}(XI\{\|X\| \leq b_n\})$, $n \geq 1$.

Remark 3.1. Under the assumptions of Theorem 3.1, we conclude from Theorem 3.1 that, for every sequence $\{X, X_n; n \geq 1\}$ of i.i.d. $\mathbf{B}$ -valued random variables,

$$\frac{S_n - \gamma_n}{b_n} \rightarrow_{\mathbb{P}} 0 \text{ if and only if } \lim_{n \rightarrow \infty} n\mathbb{P}(\|X\| > b_n) = 0,$$

$$\limsup_{n \rightarrow \infty} \mathbb{P}\left(\frac{\|S_n - \gamma_n\|}{b_n} > \lambda\right) > 0 \quad \forall \lambda > 0 \text{ if and only if } \limsup_{n \rightarrow \infty} n\mathbb{P}(\|X\| > b_n) > 0.$$

Hence

$$\frac{S_n - \gamma_n}{b_n} \not\rightarrow_{\mathbb{P}} x \quad \forall x \in \mathbf{B} \setminus \{0\}.$$

Let $0 < p \leq 2$. Then $\mathbf{B}$ is said to be of *stable type p* if

$$\sum_{n=1}^{\infty} \Theta_n v_n \text{ converges a.s. whenever } \{v_n : n \geq 1\} \subseteq \mathbf{B} \text{ with } \sum_{n=1}^{\infty} \|v_n\|^p < \infty,$$

where $\{\Theta_n; n \geq 1\}$ is a sequence of i.i.d. stable random variables each with characteristic function $\psi(t) = \exp\{-|t|^p\}$, $-\infty < t < \infty$. A remarkable characterization of stable type p Banach spaces via the WLLN was provided by Marcus and Woyczyński (1979) who showed that, for given $1 \leq p < 2$, the following two statements are equivalent:

- (i) The Banach space $\mathbf{B}$ is of stable type p .
- (ii) For every symmetric sequence $\{X, X_n; n \geq 1\}$ of i.i.d. $\mathbf{B}$ -valued variables,

$$\frac{S_n}{n^{1/p}} \rightarrow_{\mathbb{P}} 0 \text{ if and only if } \lim_{n \rightarrow \infty} n\mathbb{P}(\|X\| > n^{1/p}) = 0.$$

Combining Theorem 3.1 and the above characterization of stable type p Banach spaces, we immediately obtain the following two results.

Corollary 3.1. Let $1 \leq p < 2$ and let $\{a_n; n \geq 1\}$ be an increasing sequence of positive real numbers such that

$$\lim_{n \rightarrow \infty} a_n = \infty \text{ and } \{n^{1/p}/a_n; n \geq 1\} \text{ is a nondecreasing sequence.}$$

Let $(\mathbf{B}, \|\cdot\|)$ be a real separable Banach space such that, for every symmetric sequence $\{X, X_n; n \geq 1\}$ of i.i.d. $\mathbf{B}$ -valued random variables, (3.2) holds. Then the Banach space $\mathbf{B}$ is of stable type p .

Corollary 3.2. Let $(\mathbf{B}, \|\cdot\|)$ be a real separable Banach space. Let $1 \leq p < 2$ and let $\{b_n; n \geq 1\}$ be a sequence of positive real numbers such that

$$\left\{b_n/n^{1/p}; n \geq 1\right\} \text{ is a nondecreasing sequence.}$$

If $\mathbf{B}$ is of stable type p , then for every sequence $\{X, X_n; n \geq 1\}$ of i.i.d. $\mathbf{B}$ -valued random variables, (3.3) and (3.4) are equivalent.

Corollary 3.3. Let $\{X, X_n; n \geq 1\}$ be a sequence of i.i.d. real-valued random variables and let $\{b_n; n \geq 1\}$ be a sequence of positive real numbers such that $b_n/n^{1/p}$ is nondecreasing for some $p \in [1, 2)$. Set $S_n = \sum_{i=1}^n X_i$, $n \geq 1$. Then

$$\frac{S_n - n\mathbb{E}(XI\{|X| \leq b_n\})}{b_n} \rightarrow_{\mathbb{P}} 0 \quad \text{or} \quad \limsup_{n \rightarrow \infty} \mathbb{P}\left(\frac{|S_n - n\mathbb{E}(XI\{|X| \leq b_n\})|}{b_n} > \lambda\right) > 0 \quad \forall \lambda > 0$$

according as

$$\lim_{n \rightarrow \infty} n\mathbb{P}(|X| > b_n) = 0 \quad \text{or} \quad \limsup_{n \rightarrow \infty} n\mathbb{P}(|X| > b_n) > 0.$$

Proof. It is well known that the real line $\mathbb{R}$ is of stable type p for all $p \in [1, 2)$ and so the corollary follows immediately from Corollary 3.2. $\square$

Remark 3.2. Corollary 3.3 is an improved version of Theorem 1 (ii) of Klass and Teicher (1977) wherein $\{b_n/n; n \geq 1\}$ is nondecreasing. Theorem 1 of Klass and Teicher (1977) may be regarded as a WLLN analogue of Feller's (1946) extension of Marcinkiewicz-Zygmund SLLN (see, e.g., Chow and Teicher (1997, p. 125)) to more general norming sequences.

To prove Theorem 3.1, we use the following two preliminary lemmas. The second lemma is due to Li, Liang, and Rosalsky (2016).

Lemma 3.1. Let $\{a_n; n \geq 1\}$ and $\{b_n; n \geq 1\}$ be increasing sequences of positive real numbers satisfying (3.1). Then there exist two continuous and increasing functions $\varphi(\cdot)$ and $\psi(\cdot)$ defined on $[0, \infty)$ such that (1.1) holds and

$$\varphi(0) = \psi(0) = 0, \quad \varphi(n) = a_n, \quad \psi(n) = b_n, \quad n \geq 1. \quad (3.5)$$

Proof. Let $a_0 = b_0 = 0$. Let

$$\varphi(t) = a_{n-1} + (a_n - a_{n-1})(t - n + 1), \quad n - 1 \leq t < n, \quad n \geq 1$$

and

$$\psi(t) = b_{n-1} + (b_n - b_{n-1})(t - n + 1), \quad n - 1 \leq t < n, \quad n \geq 1.$$

Clearly, $\varphi(\cdot)$ and $\psi(\cdot)$ are two continuous and increasing functions defined on $[0, \infty)$ such that (3.5) holds. We now verify that (1.1) holds with the chosen $\varphi(\cdot)$ and $\psi(\cdot)$. Note that (3.1) implies that, for $n - 1 < t < n$ and $n \geq 1$,

$$\left(\frac{\psi(t)}{\varphi(t)}\right)' = \frac{\psi'(t)\varphi(t) - \psi(t)\varphi'(t)}{\varphi^2(t)} = \frac{a_{n-1}b_n - b_{n-1}a_n}{\varphi^2(t)} = \frac{a_{n-1}a_n \left(\frac{b_n}{a_n} - \frac{b_{n-1}}{a_{n-1}}\right)}{\varphi^2(t)} \geq 0,$$

where $b_0/a_0 \triangleq b_1/a_1$. Thus (1.1) follows. $\square$

Lemma 3.2. (Corollary 1.3 of Li, Liang, and Rosalsky (2017)) *Let $\{X, X_n; n \geq 1\}$ be a sequence of i.i.d. $\mathbf{B}$ -valued random variables. Let $\{X'_n; n \geq 1\}$ be an independent copy of $\{X_n; n \geq 1\}$. Write $S_n = \sum_{i=1}^n X_i$, $S'_n = \sum_{i=1}^n X'_i$, $n \geq 1$. Let $\{b_n; n \geq 1\}$ be an increasing sequence of positive real numbers such that $\lim_{n \rightarrow \infty} b_n = \infty$. Then we have*

$$\frac{S_n - n\mathbb{E}(XI\{\|X\| \leq b_n\})}{b_n} \rightarrow_{\mathbb{P}} 0$$

if and only if

$$\frac{S_n - S'_n}{b_n} \rightarrow_{\mathbb{P}} 0.$$

With the preliminaries accounted for, Theorem 3.1 may be proved.

Proof of Theorem 3.1 To establish the conclusion of Theorem 3.1, it suffices to show that, for every sequence $\{X, X_n; n \geq 1\}$ of i.i.d. $\mathbf{B}$ -valued random variables, the following three statements are equivalent:

$$\frac{S_n - \gamma_n}{b_n} \rightarrow_{\mathbb{P}} 0, \quad (3.6)$$

$$\lim_{n \rightarrow \infty} \mathbb{P} \left(\frac{\|S_n - \gamma_n\|}{b_n} > \lambda \right) = 0 \text{ for some constant } \lambda \in (0, \infty), \quad (3.7)$$

$$\lim_{n \rightarrow \infty} n\mathbb{P}(\|X\| > b_n) = 0. \quad (3.8)$$

Here and below $\gamma_n = n\mathbb{E}(XI\{\|X\| \leq b_n\})$, $n \geq 1$.

Let $\{X', X'_n; n \geq 1\}$ be an independent copy of $\{X, X_n; n \geq 1\}$ and set $S'_n = \sum_{i=1}^n X'_i$, $n \geq 1$. Since (3.7) obviously follows from (3.6), it suffices to establish the implications “(3.6) $\Rightarrow$ (3.8)”, “(3.8) $\Rightarrow$ (3.6)”, and “(3.7) $\Rightarrow$ (3.6)”.

It follows from (3.6) that

$$\frac{S_n - S'_n}{b_n} \rightarrow_{\mathbb{P}} 0.$$

and hence by the remarkable Lévy inequality in a Banach space setting (see, e.g., see Proposition 2.3 of Ledoux and Talagrand (1991)), we have that for every $n \geq 1$,

$$\mathbb{P} \left(\frac{\max_{1 \leq i \leq n} \|X_i - X'_i\|}{b_n} > t \right) \leq 2\mathbb{P} \left(\frac{\|S_n - S'_n\|}{b_n} > t \right) \rightarrow 0 \text{ as } n \rightarrow \infty \quad \forall t \geq 0. \quad (3.9)$$

Note that $\{X - X', X_n - X'_n; n \geq 1\}$ is a sequence of i.i.d. $\mathbf{B}$ -valued random variables. Thus (3.9) implies that

$$\mathbb{P} \left(\frac{\max_{1 \leq i \leq n} \|X_i - X'_i\|}{b_n} > t \right) = 1 - \left(1 - \mathbb{P} \left(\frac{\|X - X'\|}{b_n} > t \right) \right)^n \rightarrow 0 \text{ as } n \rightarrow \infty \quad \forall t \geq 0$$

which is equivalent to

$$n\mathbb{P} \left(\frac{\|X - X'\|}{b_n} > t \right) \rightarrow 0 \text{ as } n \rightarrow \infty \quad \forall t > 0$$

and hence (3.8) holds since

$$\{\|X'\| \leq b_n/2, \|X\| > b_n\} \subseteq \{\|X - X'\| > b_n/2\} \quad \text{and} \quad \lim_{n \rightarrow \infty} \mathbb{P}(\|X'\| \leq b_n/2) = 1$$

ensures that for all large n ,

$$\mathbb{P}(\|X\| > b_n) \leq 2\mathbb{P}(\|X - X'\| > b_n/2).$$

Thus we see that (3.6) implies (3.8).

We now show that (3.8) implies (3.6). Suppose that (3.8) holds. Set

$$\tilde{X} = \frac{X - X'}{2}, \quad \tilde{X}_n = \frac{X_n - X'_n}{2}, \quad n \geq 1.$$

Clearly,

$$\mathbb{P}(\|\tilde{X}\| > t) \leq \mathbb{P}(\|X\| > t) + \mathbb{P}(\|X'\| > t) = 2\mathbb{P}(\|X\| > t) \quad \forall t > 0.$$

Thus $\{\tilde{X}, \tilde{X}_n; n \geq 1\}$ is a sequence of i.i.d. symmetric $\mathbf{B}$ -valued random variables such that

$$\lim_{n \rightarrow \infty} n\mathbb{P}(\|\tilde{X}\| > b_n) = 0. \quad (3.10)$$

Since $\{a_n; n \geq 1\}$ and $\{b_n; n \geq 1\}$ are increasing sequences of positive real numbers satisfying (3.1), by Lemma 3.1 there exist two continuous and increasing functions $\varphi(\cdot)$ and $\psi(\cdot)$ defined on $[0, \infty)$ such that both (1.1) and (3.5) hold. Write

$$Y = \varphi\left(\psi^{-1}(\|\tilde{X}\|)\right) \frac{\tilde{X}}{\|\tilde{X}\|}, \quad Y_n = \varphi\left(\psi^{-1}(\|\tilde{X}_n\|)\right) \frac{\tilde{X}_n}{\|\tilde{X}_n\|}, \quad n \geq 1.$$

It is easy to see that

$$\mathbb{P}(\|Y\| > a_n) = \mathbb{P}\left(\left\|\varphi\left(\psi^{-1}(\|\tilde{X}\|)\right) \frac{\tilde{X}}{\|\tilde{X}\|}\right\| > \varphi(n)\right) = \mathbb{P}(\|\tilde{X}\| > b_n), \quad n \geq 1.$$

It thus follows from (3.10) that $\{Y, Y_n; n \geq 1\}$ is a sequence of i.i.d. symmetric $\mathbf{B}$ -valued random variables such that

$$\lim_{n \rightarrow \infty} n\mathbb{P}(\|Y\| > a_n) = 0$$

and hence by (3.2)

$$\frac{\sum_{i=1}^n Y_i}{a_n} \rightarrow_{\mathbb{P}} 0. \quad (3.11)$$

By Theorem 1.1 (ii) together with (3.11) and (3.10), we have that for $\epsilon > 0$,

$$\begin{aligned} \mathbb{P}\left(\left\|\sum_{i=1}^n \tilde{X}_i\right\| > \epsilon b_n\right) &\leq 4\mathbb{P}\left(\left\|\sum_{i=1}^n \varphi\left(\psi^{-1}(\|\tilde{X}_i\|)\right) \frac{\tilde{X}_i}{\|\tilde{X}_i\|}\right\| > \epsilon a_n\right) + \sum_{i=1}^n \mathbb{P}(\|\tilde{X}_i\| > b_n) \\ &= 4\mathbb{P}\left(\left\|\sum_{i=1}^n Y_i\right\| > \epsilon a_n\right) + n\mathbb{P}(\|\tilde{X}\| > b_n) \\ &\rightarrow 0 \quad \text{as } n \rightarrow \infty \end{aligned}$$

and hence

$$\frac{S_n - S'_n}{2b_n} = \frac{\sum_{i=1}^n \tilde{X}_i}{b_n} \rightarrow_{\mathbb{P}} 0.$$

We thus conclude that

$$\frac{S_n - S'_n}{b_n} \rightarrow_{\mathbb{P}} 0.$$

By Lemma 3.2, (3.6) follows.

It remains to show that (3.7) implies (3.6). Let $\{X, X_n; n \geq 1\}$ be a sequence of i.i.d. $\mathbf{B}$ -valued random variables satisfying (3.7). By the Lévy inequality in a Banach space setting, we have that for all $n \geq 1$,

$$\mathbb{P}\left(\frac{\max_{1 \leq i \leq n} \|X_i - X'_i\|}{b_n} > 2\lambda\right) \leq 2\mathbb{P}\left(\frac{\|S_n - S'_n\|}{b_n} > 2\lambda\right) \leq 4\mathbb{P}\left(\frac{\|S_n\|}{b_n} > \lambda\right) \quad \forall \lambda > 0.$$

Then it follows from (3.7) that

$$\lim_{n \rightarrow \infty} n\mathbb{P}\left(\frac{\|X - X'\|}{b_n} > 2\lambda\right) = 0; \quad \text{i.e.,} \quad \lim_{n \rightarrow \infty} n\mathbb{P}\left(\left\|\frac{X - X'}{2\lambda}\right\| > b_n\right) = 0 \quad \forall \lambda > 0.$$

That is, (3.8) holds with X replaced by symmetric random variable $(X - X')/(2\lambda)$. Since (3.6) and (3.8) are equivalent, we conclude that

$$\frac{\sum_{i=1}^n \frac{X_i - X'_i}{2\lambda}}{b_n} \rightarrow_{\mathbb{P}} 0; \quad \text{i.e.,} \quad \left(\frac{1}{2\lambda}\right) \frac{S_n - S'_n}{b_n} \rightarrow_{\mathbb{P}} 0.$$

Thus

$$\frac{S_n - S'_n}{b_n} \rightarrow_{\mathbb{P}} 0$$

which, by Lemma 3.2, implies (3.6). $\square$

Acknowledgments

The research of Deli Li was partially supported by a grant from the Natural Sciences and Engineering Research Council of Canada (grant #: RGPIN-2014-05428) and the research of Han-Ying Liang was partially supported by the National Natural Science Foundation of China (grant #: 11271286).

References

1. Chow, Y.S., and Teicher, H. 1997. *Probability Theory: Independence, Interchangeability, Martingales*, 3rd ed. Springer-Verlag, New York.
2. Feller, W. 1946. A limit theorem for random variables with infinite moments. *Amer. J. Math.* **68**:257-262.
3. Hoffmann-Jørgensen, J. 1974. Sums of independent Banach space valued random variables. *Studia Mathematica* **52**:159-186.
4. Kahane, J.-P. 1985. *Some Random Series of Functions*, 2nd ed. Heath Math. Monographs, 1968. Cambridge Univ. Press.
5. Klass, M., and Teicher, H. 1977. Iterated logarithm laws for asymmetric random variables barely with or without finite mean. *Ann. Probab.* **5**:861-874.
6. Ledoux, M., and Talagrand, M. 1991. *Probability in Banach Spaces: Isoperimetry and Processes*. Springer-Verlag, Berlin.

7. Lévy, P. 1937. *Théorie de L'addition des Variables Aléatoires*. Gauthier-Villars, Paris.
8. Li, D., Liang, H.-Y., and Rosalsky, A. 2017. A note on symmetrization procedures for the laws of large numbers. *Statist. Probab. Lett.* **121**: 136-142
9. Marcus, M. B., and Woyczyński, W. A. 1979. Stable measures and central limit theorems in spaces of stable type. *Trans. Amer. Math. Soc.* **251**:71-102.